

Economic Evaluation of Mental Health Effects of Flooding using Bayesian Networks

Tabassom Sedighi ¹, Liz Varga ², Amin Hosseinian-Far ³ and Alireza Daneshkhah ^{4,*}

¹ Centre for Environmental and Agricultural Informatics, School of Water, Energy and Environment (SWEE), Cranfield University, Cranfield MK43 0AL, UK; t.sedighi@cranfield.ac.uk (T.S.)

² Department of Civil, Environmental and Geomatic Engineering, Faculty of Engineering, UCL, London WC1E 6BT, UK; l.varga@ucl.ac.uk (L.Z.)

³ Centre for Sustainable Business Practices, University of Northampton, University Drive, NN1 5PH, Northampton, UK; amin.hosseinianfar@northampton.ac.uk (A.H.)

⁴ Research Centre for Computational Science and Mathematical Modelling, School of Computing, Electronics and Maths, Coventry University, Coventry, CV1 5FB, UK (A.D.).

* Correspondence: ac5916@coventry.ac.uk

Abstract: Appraisal of appropriate levels of investment for devising flooding mitigation and support recovery interventions is a complex and challenging task. Evaluation must account for social, political, environmental and other conditions, such as flood state expectations and local priorities. The evaluation method should be able to quickly identify evolving investment needs as the incidence and magnitude of flood events continue to grow. Quantification is essential and must consider multiple direct and indirect effects on flood related outcomes. The method proposed in this study is a Bayesian Network which may be used ex-post for evaluation, but also ex-ante for future assessment, and near real-time for reallocation of investment into interventions. The particular case we study is the effect of flood interventions upon mental health which is a gap in current investment analyses. Natural events such as floods expose people to negative mental health disorders including anxiety, distress, and post-traumatic stress disorder. Such outcomes can be mitigated or exacerbated not only by state funded interventions, but by individual and community skills and experience. Success is also dampened when vulnerable and previously exposed victims are affected. Current measures evaluate solely the effectiveness of interventions to reduce physical damage to people and assets. This paper contributes a design for a Bayesian network that exposes causal pathways and conditional probabilities between interventions and mental health outcomes as well as providing a tool which can readily indicate the level of investment needed in alternative interventions based on desired mental health outcomes.

Keywords: Bayesian network; Cost-effectiveness Intervention; Evaluation; Flood risk management; Mental health impacts; QALY

Citation: Sedighi, T.; Varga, L.; Hosseinian-Far, A.; Daneshkhah, A. Economic Evaluation of Mental Health Effects of Flooding using Bayesian Networks. *Int. J. Environ. Res. Public Health* **2021**, *1*, 0. <https://doi.org/>

Received:

Accepted:

Published:

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Copyright: © 2021 by the authors. Submitted to *Int. J. Environ. Res. Public Health* for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Natural hazards can have large societal impacts. It is estimated that they caused 7700 human fatalities and \$110 billion loss of infrastructural assets worldwide just in 2014 [1]. Out of the set of natural hazards, flooding is often regarded as the most frequently-occurring type of natural disaster with increasing risk to society (particularly, in the UK and Europe), and with the greatest impact on human [2]. Of the €150bn in reported damages caused by natural hazards in Europe in the period of 1999 to 2009, over one-third of damages (i.e. €50bn) were due to flooding. Furthermore, annual flood losses are expected to increase five-fold by 2050 and nearly 17-fold by 2080 in Europe, drawing attention to the urgency for cities in Europe to construct resilience against flooding [3].

Similar to other natural disasters, when flooding occurs, it creates significant damage to homes, communities, businesses, public services, etc. Residential properties




Figure 1. There are direct and indirect impacts of flood damages that are not easily be quantified in monetary terms [4]

usually suffer the greatest proportion of flood damage, with 25% of total damage e.g., £320 million cost incurred by 10,465 properties due to flooding [5]. Therefore, flood risk management is a disaster administration priority for European countries, particularly the UK.

It has been argued that flood risk management is usually measured as direct property and infrastructure losses, since these are the most important input for cost–benefit analysis that guide the government bodies to invest in flood risk management strategies [6]. However, the impacts of flooding on urban populations are multi-faceted and wide-ranging. It is well-known that floods also have enormous impacts on people, both directly and indirectly (see Figure 1). Distinctions must be made between direct/indirect and tangible/intangible flood damages. Direct damage becomes immediately visible in the affected areas due to close physical contact with floodwater, while indirect damage emerges with a time delay and/or outside the area affected by floods [4].

The most apparent intangible impact of flooding is on human health. Direct intangible damage is a primary loss, which manifests as physical injury or even loss of life. Indirect health impacts are mental health disorders, which are caused by the experience of being flooded, or being impacted during the restoration process. Estimating flooding impacts will be provide valuable insights for decision making and risk mitigation, policy-making, civil protection, emergency alertness and response, insurance and reinsurance, damage estimation practice/research, etc. [7].

On account of this, an comprehensive societal cost–benefit assessment must take into account intangible losses caused by floods, such as psychological disorders or anxiety [8], as well as tangible losses. Due to anticipated complications of converting intangible values, such losses are generally ignored in risk assessments [9]. Thus, economic evaluation of the convincing levels of investment which should be made into interventions to mitigate the flood risk, and support recovery from floods is very challenging.

In this paper, the primary focus is on the evaluation of flood impacts on human health, particularly mental health [10]. Flood impact assessment is a key component of the practice of flood risk management. Flood risk is defined in the European Flood Directive as “the combination of the probability of a flood event and of the potential adverse impacts on human health, the environment, cultural heritage and economic

66 activity associated with a flood event” [11]. Flood damage estimates are therefore, useful
67 at all the stages of what is known as the flood mitigation cycle.

68 It is thus crucial to embrace social, political, environmental and other conditions,
69 such as flood likelihood and local priorities, into the comprehensive evaluation. The eval-
70 uation method must be also able to swiftly determine changed investment requirements
71 as the incidence and magnitude of flood events continue to grow. This quantification is
72 essential and must examine various direct and indirect flood impacts on flood related
73 outcomes in a probabilistic manner.

74 In this paper, we illustrate how the Bayesian network (BN) probabilistic method
75 can be used efficiently for ex-post economic evaluation, as well as ex-ante for future
76 assessment, and indeed near real-time for reallocation of investment into interventions.

77 In general, BNs provide a robust and flexible analytic approach to the challenge
78 of complex health datasets, which pose specific computational challenges because of
79 missing data, large or small size of data, complexity (of relationships not only between
80 variables but also in the datasets themselves), changing populations, and nonlinear
81 relationships between exposures and outcomes [12]. Unlike the regression-based models
82 or multivariate copula models [13]—the BNs historically most commonly used in clinical
83 risk prediction analysis and risk stratification [14] in medicine. They provide compact
84 and instinctive graphical representations that can be used to conduct causal reasoning
85 and risk prediction analysis. Furthermore, the cause and effect statements can be readily
86 exploited in BN networks to reduce the computational time and cost of. This can be
87 considered as another important advantage of this modelling approach in comparison to
88 the conventional approaches, such as joint probability distribution, which only encodes
89 the values of the outcomes of interest, given the input variables. Therefore, Bayesian
90 networks offer a compact tool for dealing with the uncertainty and complexity of a
91 system. In this study, the benefits, efficiency and limitations of the BN-based evaluation
92 method will be studied by examining the effect of flood interventions upon mental
93 health which is a gap in current investment analyses.

94 In order to construct the proposed probabilistic methods, we need to have a holistic
95 overview of the relationship between flood events, their aftermath, and population
96 well-being and risk factors causing psychological disorders. The psychological health
97 impacts of flooding, and their relationship with flooding and other risk factor will be
98 briefly discussed in the next section. Estimating the cost of flooding on human health,
99 in particular on human mental health, is very challenging but essential in order that
100 investment into interventions can be evaluated against reduced mental health impacts.
101 It is essential to use metrics/methods to monetize mental health impacts.

102 In order to reduce the damages caused by flood events on the community and
103 people, environmental agencies are using various interventions each with different
104 outcomes, efficiencies and costs. Any combination of interventions results in different
105 value for money, with multiple conditional dependencies between interventions, choices
106 of implementation and their contexts. This study provides an efficient construction
107 for a probabilistic BN that displays causal pathways and their probabilities between
108 interventions and mental health outcomes as well as providing a tool which can readily
109 indicate the level of investment needed in alternative interventions based on anticipated
110 mental health outcomes.

111 2. Psychological Impacts of Flooding

112 The psychological impacts of flooding can be very significant and long-lasting.
113 Difficulties in evaluating mental health impacts of flooding arise because accurate
114 diagnosis of any condition is not straightforward, and mental health impacts are often
115 under-reported, and can be overlooked in comparison to the physical health impacts.

116 There are some studies evaluating the impact of flooding on mental health. In
117 one of the earlier studies, [15] conducted a study to evaluate the psychological impacts
118 attributed to a severe flooding in Kentucky, US in 1984. The findings indicated that

119 the flood exposure had psychological impacts on the population and impacts included
120 depression and anxiety. [16] conducted a similar study with a group of participants
121 from a flood affected population in the town of Lewes in the UK, to evaluate both the
122 physical and mental health effects of the flooding in the area in the year 2000. The
123 study findings identified a high correlation between flood exposure and psychological
124 distress. Such physical and psychological consequences denote people's vulnerabilities
125 as they interact with nature [17]. Tapsell et al. go on to assert that quantification of
126 natural disaster impacts on population health is an intricate task due to the delay in
127 receiving feedback from the population. Nevertheless, they conducted a similar piece
128 of research on the impacts on flooding of 1998 in large parts of England and Wales.
129 Their longitudinal study took place over a period of four and half years, evaluated
130 both physical and psychological impacts. The top four psychological impacts in the
131 few weeks or months after the flood were claimed to be 'Anxiety', 'Increased Stress
132 Levels', 'Sleeping Problems', and 'Mild Depression'. Nonetheless, the order by which
133 these health effects were reported varied from one geographical area to another.

134 The UK and England in particular are prone to flooding. In 2005, a severe flood hit
135 Carlisle, UK, and many homes were affected. Carroll et al. [18] conducted qualitative
136 research to evaluate the psychological impact of this specific flood and to evaluate the
137 impacts of disasters and how they could inform policies. They concluded that the main
138 psychological impacts are anxiety, stress and post-traumatic stress disorder (PTSD).
139 Another study reported specifically that females were psychologically more vulnerable
140 than males in the event of flooding [19].

141 Research in other regions provide similar findings. Vietnam is also susceptible to
142 natural disasters and specifically to flooding. Bich et al. [20] highlight that controlling
143 communes significantly reduces psychological impacts when flooding occurs. There
144 are different strategies to mitigate the impact of flooding from low impact development
145 technologies [21], relocation [8], to forestation [22]. It was also reported that relocation
146 during flood recovery, as an intervention, is correlated with 600% increase in mental
147 health symptoms [8].

148 Zhong et al. [23] provide a better understanding of what is currently known re-
149 garding the long-term health impacts of flooding and the factors that may influence
150 health outcomes (including psychological health) by conducting a systematic mapping.
151 Their findings indicate that 68% of these studies focused on psychological impacts of
152 flooding, whereas only 16% of these studies evaluated the physical effects following
153 exposure to flooding. They have underlined that future research needs to quantify the
154 long-term health impacts of flooding and identify their major determinants using some
155 novel quantitative tools. These tools should be able to quantify the influence of multiple
156 social interventions, such as flood management, on long-term health outcomes, and also
157 identify the most influencing factors affecting the psychological and physical impacts.

158 3. Cost Estimation of Flooding

159 Estimating the cost of flooding on human health including mental health is ex-
160 tremely challenging. The study by [24] reports the best indicative estimates for the
161 loss of life and health for the 2015 to 2016 UK winter floods (i.e. £43m, within a range
162 of (£32m, £54m)). The best estimate of loss of life and health impacts is calculated as
163 "surrogate cost of fatalities" plus "surrogate cost of health impacts", where surrogate
164 cost of fatalities (£5m) is measured as the number of fatalities due to flooding times
165 'average value of prevention of fatality'. The "surrogate cost for health impacts" (£38m)
166 is calculated as 'cost per household' times 'number of households affected'. The first
167 term (cost per household) is defined as household willingness to pay per year to avoid
168 health impacts of extreme flood events, times, discount factor in the year, and the second
169 term (number of properties affected) is measured as the 'number of residential properties
170 flooded' times 'number of households likely to have health affects'.

171 Most studies focused on direct impacts. The common types of health metrics used
172 are: death; hospital admissions and out-patients visits; cases of acute morbidity or
173 injuries; and mental disorders or reduction in well-being [25].

174 However, loss of life or number of injured are commonly used to measure the health
175 burden associated with any natural disaster, and therefore, the impact of flooding on
176 individuals' mental health is often overlooked. In order to monetize health impacts
177 in the flooding context, the following should be considered: Healthcare resource use;
178 Productivity loss; Dis-utility from suffering or life-shortening.

179 The monetary value of the latter component is typically evaluated by wealth-health
180 trade-offs that the affected people reveal in surrogate markets or can be implemented
181 through multiple choice experiments. The monetary value of dis-utility associated with
182 an adverse health outcome is thus attributed to the willingness to pay (WTP) to avert
183 outcomes or, when considering mortality risk, the value of a statistical life that is derived
184 from individuals' aggregated WTP for a small change in survival probabilities [25]. In
185 the studies that used loss of life numbers to quantify health impacts, only a few of them
186 applied a monetary value to this outcome by multiplying it with a value of statistical
187 life. This is not surprising, given that monetizing death is less useful for descriptive
188 studies that are investigating trends in effects, or for studies reporting results from
189 population-based surveys.

190 Matsushima et al. [26] valued WTP to avoid mental damages from flooding using
191 an option value approach, in order to address potential strategic bias that would lead to
192 an over-valuation of WTP. The WTP was also reported in [27] to estimate the willingness
193 to contribute in labour, in order to circumvent the fact that most individuals would not be
194 able to afford any financial payment. They have also concluded that flood damage was
195 estimated on average to represent about 20% of households' annual income. However,
196 it was not possible to solve the welfare loss from morbidity and well-being reduction
197 from the welfare loss due to damages to assets. Poor households were found to be more
198 vulnerable to flooding as the associated damages made up a significantly larger
199 portion of their annual income. Households heavily dependent on agricultural activities
200 were also found to be more vulnerable.

201 The UK Environment Agency (EA) has recently studied the new economic costs
202 for the mental health impacts of flooding by analysing the data provided by Public
203 Health England (PHE). It was illustrated that the mental health prevalence of people
204 disrupted or affected by flooding is considerably higher than the unaffected groups,
205 over 12 and 24 months periods. The findings of the study are comparable to the results
206 from the flooding occurred in 2007. It was also reported that the chance of any type of
207 mental health outcomes will increase by the flood severity (or depth of flood) among the
208 affected population. The same study confirms that WTP could be a very useful metric to
209 evaluate the social cost of the flood impact, however, it cannot be used to include the
210 actual cost of the mental health outcomes to the economy.

211 A study commissioned by Defra suggested that households were, on average,
212 willing to pay £200 per year (2004 prices) to avoid the negative health impacts of flooding
213 (e.g. for events occurring less frequently than 1 in 75 years) [28]. Defra's climate change
214 risk assessment report [29] considers the costs of treating a case of mild depression
215 following a flood event as £970 (2010 prices), which can be used as an indicator of mental
216 health impacts. It should be noted that these monetary values are normally used as
217 predictions in policy assessments to allocate resources to protect against an abstract
218 individuals' loss of life or suffering from harm. They were not designed to include
219 post-event analysis. Without any official post-event values, however, these values were
220 used as a surrogate in both the 2007 and 2013 to 2014 'cost of flood reports' to provide an
221 indicative sum for loss of life and health impacts. Nevertheless, the above research and
222 other studies conducted by PHE intended to better understand the health impacts of
223 flooding and these efforts have resulted in some changes in the 2013 to 2014 methodology
224 for estimating cost of flood. More research is urgently required to estimate the cost of

225 treating cases of anxiety, depression and PTSD, using the existing and other relevant
226 data. The factors affecting the cost of mental health outcomes are:

- 227 • Knowledge of each outcome (or condition),
- 228 • Prevalence of these outcomes,
- 229 • Presence of known treatment plans,
- 230 • Duration of any treatment,
- 231 • Likely impact of the outcome over the short term in terms of days of work lost.

232 The quantification of the benefits of flood risk prevention measures is still an
233 unresolved challenge in disaster management research works. In particular, there is
234 no clear flood risk management to quantify the effect of interventions in reducing the
235 flooding impacts on people including the effects on the affected population's mental
236 health. The most widely adopted framework in flood risk reduction is represented as the
237 calculation of the expected damages as a function of flood hazard, physical vulnerability
238 and exposure [30]. According to this framework, flood hazard is characterized by specific
239 return periods – an estimate of the likelihood of the flood. Moreover, together with
240 the vulnerability, it is usually expressed as an index, while the exposure is expressed
241 with the unit(s) of measurement of the elements at risk, in physical or monetary terms.
242 However, floods can impact socio-ecological systems in various forms, and therefore this
243 framework is limited to assess damages to constructed infrastructure only. Furthermore,
244 there have been a few other attempts to provide such a holistic risk assessment (see [31,
245 32]), yet, these methods primarily focus on assessing direct tangible costs, since there is
246 only enough relevant information to justify decisions regarding structural risk reduction
247 measures. The main challenge with traditional frameworks is that they neglect the
248 fact that the magnitude of flooding costs is determined by the adaptive behaviour of
249 communities to absorb the flood hazards. It is obvious that the human dimension of
250 vulnerability must be addressed as one of the main elements of the flood risk. The
251 human aspect of vulnerability relates to the ability to cope with the hazard after a flood
252 and the capacity to adapt to the flood hazard before the event [33]. More recently,
253 emphasis has shifted from just being prepared, informed, and minimising the dimension
254 of vulnerability, to strategic proactive planning and management. There are two main
255 reasons for this shift in recent flood risk management:

- 256 1. The uncertainty of flood occurrence has noticeably increased due to intensified
257 climate change; and
- 258 2. The consequences of flooding considerably depend on the behaviour of the affected
259 people and their capability to adapt.

260 4. Mitigating the Impact of Flood Health Damages

261 There are various interventions that reduce the damages caused by flood events
262 on communities, local environmental agencies, with different outcomes, efficiencies
263 and costs. For instance, an intervention can be to use an early warning system (EWS)
264 to reduce the amount of direct tangible costs (e.g., people can move transportable
265 properties outside of the exposed area when the flood hazard is anticipated). The aim of
266 a flood warning system is to provide useful information, for instance, by issuing alerts
267 or activating the required protection measures with a view to improve decision making
268 and action. The connections and feedback between hydrological and social spheres of
269 early warning systems are key elements of a successful flood mitigation.

270 The behaviour of the public and first responders during flood situations, are deter-
271 mined by their preparedness, and are heavily influenced by numerous behavioural traits
272 such as perceived benefits of protection measures, risk awareness, or even denial of the
273 effects that might occur. In the UK, the Environment Agency (EA) has an important
274 role in warning citizens about the risk of flooding with a view to reduce the impact of
275 flooding from rivers and the sea as well as pluvial floods.

276 In November 2009, Cumbria experienced devastating flooding in its different re-
277 gions due to the heaviest rainfall ever recorded in the UK [34]. Following this, the

EA carried out qualitative and quantitative research to evaluate the impact of the EA's flood intervention methods, including early warning systems, partnership work, and on-ground assistance. These research works also highlight opportunities to improve the EA's ability to respond to future floods. The affected residents received warnings in the various forms: including EA Flood-line Warnings Direct, people own observation of the local area, warnings on weather forecasts, warnings from neighbours, friends, and/or family, the Flood-line, and warnings by the emergency services [35]. They found that early warning systems themselves could add to stress. Also, most people found Flood Action Groups very helpful in protecting their homes against flooding. However, they valued the idea of making a flood action plan, though such flood action planning was not yet widespread.

Another intervention to reduce mental health harm is to relocate people away from the affected regions as soon as possible, and to support them during and after a flood. This also has rebound effects.

A probabilistic method is needed to consider all sources of uncertainty that may influence an intervention in a particular context in order to evaluate its value for money. There are complex relationships between flood events, their aftermath, population well-being and risk factors causing people's health deterioration and/or psychological disorders [36].

To understand potential benefits or drawbacks of any intervention for reducing the damaging impacts on people's health, and particularly their mental health, needs to take into consideration the nature of the hazard, the vulnerability of the community and its exposure. Figure 2 illustrates a conceptual model of a customised version of the risk framework considering the impacts that EWS may have on people [33]. In this framework, Hazard refers to the potential occurrence of flood which may cause loss of life, injury, or other health impacts, as well as damage and loss of property, infrastructure, livelihoods, service provision, and environmental resources.




Figure 2. Customised application of the risk framework by including early warning system (some information derived from the original framework developed by [33])

304

305 5. Flooding and Health Risk Factors: Modelling approaches

306 There are currently several statistical methods to explore the relationship between
 307 flooding and the health risk factors discussed above. For instance, a multivariable
 308 logistic regression model was proposed by [37] to model individuals' revealed changes
 309 in mental health outcomes between year one and year two after flooding, by considering

310 some of the above-mentioned factors. A similar method (logistic regression analysis)
 311 was used to select the risk factors and to predict the flooding victims' mental health
 312 states [38]. Applications of the multivariate regression-based methods are very limited.
 313 Their performance is hugely dependent on the size of the dataset, and can hardly be
 314 used to efficiently model the complex relationships between flood events, their aftermath
 315 impacts, and risk factors causing people's health deterioration and/or psychological
 316 disorders. In addition, they are not useful in assessing risks in complex systems and
 317 scenarios of 'decision making under uncertainty' to optimise cost-effective decisions.

318 Alternatively, probabilistic methods, particularly BNs have become an increasingly
 319 popular method for modelling uncertain and complex systems [39] and are considered
 320 as a powerful tool for presenting knowledge and interpreting insights from available
 321 data [40]. Applications of BN methods are found in a growing number of studies, and
 322 disciplines [41]. BNs are particularly useful for evaluation due to their capability of
 323 classification based on observations. BNs have been also widely used in environmental
 324 management contexts and are appropriate for decision making under uncertainty [42,43].
 325 Moreover, unsupervised learning from a dataset can be performed using a BN by
 326 adopting the learning algorithm to find both structure and conditional probabilities. This
 327 means the evaluator does not need to know how to create a BN, although it is possible
 328 to aid the learning algorithm with a prior knowledge about relations and probabilities.
 329 Dealing with uncertainty when evaluating policy is a challenge that can be addressed
 330 using BNs, since uncertain probabilities of variables may be safely ignored to get to the
 331 desired probabilistic quantity of a random variable. Furthermore, BNs engage directly
 332 with subjective data in a transparent way. Hence, the method could be considered as a
 333 tool to explore beliefs, evidence and their logical implications, than as a means to 'prove'
 334 something in somewhat absolute sense. They, therefore, are useful in producing the
 335 balanced judgements required for evaluation in a Value for Money context. Additionally,
 336 BNs can be used privately to structure and inform the evaluator's understanding, or
 337 publicly in a participatory process to stimulate and challenge collective views [41].
 338 Finally, BNs are user-friendly, and practical, and can present intuitively and graphically
 339 the 'story' behind a finding.

340 6. Evaluation method: Probabilistic Graphical Models

341 Bayesian network (BN) is a mathematical model that graphically and numerically
 342 represents the probabilistic relationships between random variables through the Bayes
 343 theorem. This technique is becoming popular to aid in decision-making in several
 344 domains due to the evolution of the computational capacity that makes possible the
 345 calculation of complex BN [44]. Applications of BN methods are found in a growing
 346 number of disciplines and policies [14,41,45,46].

In the BN, as a probabilistic graphical model which is used to represent knowledge
 about an uncertain domain [44], each random variable is represented by a node in the
 BN. The BN, \mathcal{B} , is a directed acyclic graph (DAG) that represents a joint probability
 distribution over a set of random variables $\mathbf{X} = (X_1, X_2, \dots, X_n)$. The network is defined
 by the pair $\mathcal{B} = \{\mathcal{G}, \theta\}$, where $\mathcal{G} = (\mathbf{X}, E)$ is a DAG with nodes \mathbf{X} representing random
 variables and edges E representing the direct dependencies between these variables. θ is
 the set of probability functions (i.e., node probability table) which contains the parameter
 $\theta_{x_i|pa_i} = P_{\mathcal{B}}(x_i|pa_i)$ for each x_i in X_i conditioned by the parent set of x_i , denoted by pa_i ,
 as the set of parameters of X_i in \mathcal{G} . The joint probability distribution defined by \mathcal{B} over
 \mathbf{X} is given in Eq. (1):

$$P_{\mathcal{B}}(X_1, \dots, X_n | \theta) = \prod_{i=1}^n P_{\mathcal{B}}(X_i | pa_i) = \prod_{i=1}^n \theta_{x_i|pa_i} \quad (1)$$

347 A simple example of a BN is illustrated in Figure 3, where the probability of a person
 348 having cancer can be computed in terms of "Relatives had cancer" (Y_1) and the person is
 349 smoking or not (denoted by Y_2).




Figure 3. A simple BN model indicating the inter-dependencies between lung cancer classifier and the affecting risk factors (adopted from [47]).

350 As it can be observed, a conditional probability table (CPT) is attached to each
 351 node. The CPT on each node is associated with the conditional probability distribution,
 352 as given in Eq. (1). **The CPTs (or conditional probabilities) can be estimated from the**
 353 **observed data or expert opinions [14,48].** A link (or ‘edge’) between two nodes represents
 354 a probabilistic dependency between the linked nodes. The links are shown with an arrow
 355 pointing from the causal node(s) (Y_1, Y_2 in Figure 3) to the effect node (X : Lung cancer
 356 in Figure 3). There must not be any directed cycles: one cannot return to a node simply
 357 by following a series of directed links. Nodes without a child node are called leaf nodes,
 358 nodes without a parent node are called root nodes (Y_1, Y_2), and nodes with parent and
 359 child nodes are called intermediate nodes. In other words, a BN represents dependence
 360 and conditional independence relationships among the nodes using joint probability
 361 distributions, with an ability to incorporate human oriented qualitative inputs. The
 362 method is well established for representing cause-effect relationships.

363 **BN learning consists of two general steps: (i) Finding DAG, which illustrates the**
 364 **inter dependency between the variables/nodes and (ii) Finding CPT for each node given**
 365 **the values of its parents on the learned DAG. Finding the best DAG is the crucial step**
 366 **in BN design. Construction of a graph to describe a BN is commonly achieved based**
 367 **on probabilistic methods, which utilise databases of records [48], such as the search**
 368 **and score approach. In this approach, a search through the space of possible DAGs is**
 369 **performed to find the best DAG. The number of DAGs, $f(p)$, as a function of the number**
 370 **of nodes, p , grows exponentially with p [49].**

371 In this paper, BN will be used to evaluate the effect of flood interventions upon
 372 mental health to explore and display causal and complex relationships between key
 373 factors and final outcomes in a straight-forward and understandable manner. The
 374 proposed BN is also used to calculate the effectiveness of the interventions, and the
 375 uncertainties associated with these causal relationships, which will be discussed in the
 376 next section. **Due to the lack of data, the proposed BN in this study was learned based**
 377 **on expert judgments (including experts from EA and Public Health of England (PHE)),**
 378 **and narrative in the relevant literature (as discussed in the next section). However, this**
 379 **approach will effectively work with data from a variety of sources, and handles a mix**
 380 **of subjective and objective data that can be incorporated with variables from different**
 381 **contexts [14,48].** Moreover, BN is a reasonable supplement to traditional statistical
 382 methods, since traditional statistical methods were unable to update complex system in
 383 the light of new information, while BNs can update the system when new evidence is
 384 added during analysis. The proposed BN developed an understanding of the effect of
 385 flood interventions, and the risk factors associated with higher impact on mental health
 386 outcomes.

387 To construct a BN for evaluation of the effect of flood interventions upon mental
 388 health, the following steps need to be performed:

- 389 1. BN structure learning: There are a number of risk factors related to the flood inter-
 390 ventions upon mental health including healthcare resources, flood management
 391 practices, existing mental disorders and many more which will be considered as
 392 input and mediate nodes to the proposed BN model. The level of effectiveness
 393 between these nodes and causal relationships between them are presented by edges,

- 394 which can be elicited from the domain experts, and available data to construct the
395 BN structure.
- 396 2. Parameter learning: prior probabilities assigned to root nodes and conditional
397 probabilities for dependent (leaf) nodes are elicited from the experts domain and
398 existing information. In the BN, the state of some nodes could be influenced by their
399 prior states, or affect other nodes. The probabilities of these nodes are determined
400 before propagating evidence to the model [50,51].
 - 401 3. Outcomes of BN (Posterior probability learning): The final step in BN is to run the
402 model at agreed intervals. As new information is added to the model, the current
403 priors/states will be updated using the Bayesian paradigm in a very efficient way.

404 It is also straightforward to use the BN to identify which variables have the largest
405 influence on the final outcomes of the network. A unique feature of BNs is the ability to
406 back propagate the model's conditional probabilities through the model structure. This
407 means that we can test how to achieve desired outcomes by identifying the most likely
408 combination of risk factors.

409 The BN model can be used to develop an effective and efficient decision support tool.
410 In the next section, the BN-based decision support tool will be developed to evaluate the
411 cost-effectiveness (in the monetary value) of various flood interventions upon mental
412 health in the present of different uncertainties and under certain constraints.

413 7. Results

414 7.1. Using Bayesian network to evaluate the effect of flood interventions upon mental health

415 In this section, we evaluate the impact of the flood interventions on the mental
416 health of the affected people by flooding using a BN trained by combination of the data
417 extracted from a narrative in the relevant literature from the published reports and expert
418 judgments (including experts from EA and PHE). However, it is very straightforward to
419 train a BN based on combination of the heterogeneous data collected from the surveys
420 and other methods [14].

421 We first need to learn the BN for a subset of the risk factors selected in relation to
422 the flood intervention upon mental health, including prevalence of probable depression
423 in people who have been flooded (Flood), loss of sentimental items (LSOI), prevalence
424 of loss of sentimental item as secondary stressor in those exposed to flooding (PPD), less
425 severe depression (Lsever), and more severe depression (Msever).

426 It is usually recommended that the BN structure and model parameters should
427 be learned from the combination of data and expert judgments [48]. However, Vepa
428 et al [14] argue that the best BN structure learned from data only, and by employing
429 various score-based or constraints-based methods [49], would not result in the model
430 favoured by the domain experts. As a result, the BN structure illustrated in Figure 4 is
431 learned based on the expert opinions only (as suggested in [14,48]). The learned BN for
432 the selected risk factors was validated by the domain expert (Economic Evidence expert
433 from the English Environment Agency). It should be noted that "CU" and "CQALY" in
434 the learned BN shown in Figure 4 stand for "change in utility" and "change in QALY",
435 respectively, which will be discussed later in this section.

436 In the next step, we need to estimate or determine the CPTs. Due to the lack
437 of data, the conditional probabilities of each node of the BN shown in Figure 4, are
438 determined using the expert opinions [52] and the information extracted from the
439 narrative of literature. Table 1 shows the summary of the probabilities of each node,
440 illustrated in Figure 4 and the source of information used to determine these probabilities.
441 For instance, the probability of LOSI is reported to be 62% based on elicited opinions
442 from the EA and PHE experts, while the prevalence of probable depression in people
443 who have been flooded (denoted by "Flood") is determined to be 20.1% [53] and the
444 English National Study of Flooding and Health (NSFH, 2020), which is available at
445 <https://bit.ly/3eXiKwt>. The probability of PPD was determined to be 18.6% [53]. The




Figure 4. Static Bayesian network to evaluate the effect of flood interventions upon mental health where LOSI indicates the loss of sentimental items, PPD indicates the prevalence of probable depression, Msever and Lsever indicate the more severe depression and less severe depression respectively, CU indicates the change in utility and CQALY indicates the change in QALY.

446 probabilities of Lsever and Msever are respectively determined to be 48.3% and 21.1%
 447 (NSFH, 2020).

Table 1: The elicited probabilities and corresponding source of data, for each node of BN illustrated in Figure 4.

Input parameter (node)	probability	Source of data
Flood	20.1%	[53] (pp. 8)
LOSI	62%	domain experts' opinions
PPD	18.6%	[53] (pp. 15)
Lsever	48.3%	(NSFH, 2020)
Msever	21.1%	(NSFH, 2020)

448 Following [54], three mental in this study: Remission, less severe depression
 449 (Lsever), and increased or more severe depression (Msever). The utility value of being in
 450 remission from depression was suggested to be 0.85, while having less severe depression
 451 is 0.60 and more severe depression is 0.42. For the sake of simplicity at this stage, we
 452 assume these utility states are monitored over one year, and that remission from de-
 453 pression is equivalent to not having depression. Moreover, there could be some overlap
 454 between the two states of Lsever and Msever, which then need to compute the change in
 455 utility as illustrated in Table 2. It should be noted the mean values reported in Table 2 are
 456 computed as the meas of suggested Beta distribution (denoted by $Be(\alpha, \beta)$ in the fourth
 457 column). These Beta distributions can be used to determine the cost-effectiveness inter-
 458 vention by optimising the Expected Value of Perfect Information measures [55,56], which
 459 is beyond the scope of this article and will be considered as the further development of
 460 this study.

Table 2: The states of mental health and their corresponding utility values, as suggested in [54].

Input parameter (health states)	Mean value	Change in utility	Probability distribution	Source of data
Remission	0.85		$Be(923, 163)$	[54]
Lsever	0.60	$(0.85 - 0.6) = 0.25$	$Be(182, 122)$	[54]
Msever	0.42	$(0.85 - 0.42) = 0.43$	$Be(54, 75)$	[54]

461 Next, the proposed BN presented in Figure 4 computes the change in QALY
 462 (CQALY) caused by loss of sentimental items (Table 3). A QALY is a measure that
 463 combines health-related quality of life and length of life into a single measure of health
 464 gain. The National Institute for Health and Clinical Excellence (NICE) provides the cost-
 465 effectiveness threshold range, which is between £20000 and £30000 per QALY [57,58].

Table 3: The change in QALY outcomes due to an intervention taken by the EA.

Health state	Before intervention	After intervention	The difference	CQALY outcomes
Msever	0.055	0.033	0.022	$0.022 \times £20000 = £440$
Lsever	0.062	0.038	0.024	$0.024 \times £20000 = £480$

466 Let assume before taking an intervention (e.g., using flood early warning system
 467 by the local EA to inform the people in advance about flood hazard) that could lead to
 468 an individual losing their sentimental item, the changes in QALY for two mental health
 469 states (i.e., Msever and Lsever) are computed from the learned BN illustrated in Figure 4
 470 as follows:

- 471 • For Msever: CQALY=0.055
- 472 • For Lsever: CQALY=0.062

473 The above CQALYs are computed based on the mean values suggested for the mental
 474 states as given in Table 2.

475 The QALY values, if the intervention was decided to be taken by the local EA prior
 476 to the flooding, will be computed (from the BN) as follows:

- 477 • For Msever: CQALY=0.033.
- 478 • For Lsever: CQALY=0.038.

479 The differences that the intervention will make for the mentioned mental health states
 480 are given by

- 481 • For Msever: $0.055 - 0.033 = 0.022$;
- 482 • For Lsever: $0.062 - 0.038 = 0.024$.

483 Finally by multiplying these changes in QALY by the lowest point of NICE's QALY
 484 cost-effectiveness threshold (e.g. £20,000), we can evaluate the cost-effectiveness of the
 485 suggested intervention on reducing the impact of the mental health due to losing of
 486 sentimental items, as:

- 487 • For Msever: $0.022 \times £20,000 = £440$.
- 488 • For Lsever: $0.024 \times £20,000 = £480$.

489 This suggests that using flood early warning system by the local EA to inform the
 490 people could save at least £480 to ensure that an individual will not suffer the less severe
 491 depression due to losing their sentimental items in the flooding events.

492 It should be noted that the early warning system could itself create further stress.
 493 An alternative way would be to relocate people away from the affected regions as soon as
 494 possible, and to support them during and after a flood. Although, any of these strategies

495 or their combinations could affect the flooded people's mental health, with each strategy
496 imposing varying benefits and costs. The method proposed above can provide us with
497 an effective cost-benefit analysis approach in comparing the suggested interventions,
498 by taking into account the complex relationships between flood events, their aftermath,
499 population wellbeing and risk factors causing people's health deterioration and/or
500 psychological disorders, and costs and benefits of the interventions.

501 8. Conclusions

502 BNs have been written to evaluate (ex-post) the effect of different factors on out-
503 comes, in contexts other than flooding. For instance, a BN has represented the in-
504 teractions of indoor climate factors on the mental performance of office workers, to
505 demonstrate that investment in improved thermal conditions is economically justified in
506 most cases with different building designs [44]. This paper is interesting because it is
507 not an evaluation of a single case, but attempts to gain insights across a diversity of past
508 cases. Clearly, if a BN accurately reflects the conditional probabilities of past cases, it can
509 be used for assessment (ex-ante) quantification of forthcoming designs of buildings, but
510 there are considerations to be addressed.

511 A BN used for evaluation, based on retrospective data and expert opinion, would
512 need to change to represent the future scenario being assessment. The scenario may
513 introduce new variables (and associated changes to prior probabilities) for example to
514 represent how the future complexity of the system will work. The scenario may require
515 changes to the values of existing variables, for example, to reflect future efficiencies
516 or effects of different ways of organizing. In fact, a number of BNs may be developed
517 to examine alternative viable futures. Every BN will have a set of unique conditional
518 probabilities which will assess the future conditions or scenarios. Across all BNs a range
519 of potential outcomes will ensue and will provide an indication of future outcomes.
520 However, scenarios are speculative and indeed deterministic [59], but if they extend cur-
521 rent representations of factors upon outcomes, they may be argued as rational extensions
522 of current understanding. In general, a BN-based modelling approach would enable us
523 to compute the costs and benefits, based on multiple causal factors including individual
524 risk factors or interventions on the system, by taking into consideration uncertainty
525 in the input parameters. The derived estimated outcomes could be presented in the
526 form of probabilities if appropriate probability distribution could be considered on the
527 input parameters (see Section 7.1). It should be noted that the BN similar to many other
528 Machine learning methods is a data-driven approach. As a result, the derived results
529 using BN are not generalisable and are fully dependent on the collected data and this
530 assumption that the data sources are accurate. We identify this as a limitation of the
531 proposed approach.

532 The health economic evaluation methodology proposed in this study, to explore
533 the level of investment required in alternative interventions based on desired mental
534 health outcome, could be developed further by computing the value of information (VoI)
535 analysis [55] (e.g., expected value of perfect information or EVPI). It should be desired
536 to compare the cost-effectiveness of the suggested interventions for the flood morbidity
537 related mental health depression from both the healthcare perspective and a societal
538 perspective, using the proposed evaluation method proposed in this study and a VoI
539 analysis, which estimates the expected value of eliminating the uncertainty surrounding
540 cost-effectiveness estimates, for both perspectives. Furthermore, the concept of expected
541 value of perfect information, which is a particular measure of VoI analysis, can be used
542 to examine probabilistic sensitivity analysis for the discussed cost-effectiveness problem
543 (see [55,56]).

544 It would be also interesting to explore further the price or threshold that a healthcare
545 decision maker or policymaker would be willing to pay to have perfect information
546 regarding all factors that influence which intervention choice is preferred as the result
547 of a cost-effectiveness analysis. However, this could be answered by VoI analysis as

548 the value (in money terms) after removing all uncertainty from such an analysis, but it
 549 requires more research and data to investigate the effectiveness of the proposed method
 550 in this regard.

Author Contributions: T.S., L.V. A.H., and A.D. contributed to the study design. T.S., L.V., A.H. and A.D. contributed to data collection. T.S. and A.D. contributed to data analysis. T.S., L.V., A.H. and A.D. contributed to original draft preparation, review and editing. All authors have read and agreed to the published version of the manuscript.

Funding: “This research received no external funding”.

Institutional Review Board Statement: “Not applicable” .

Data Availability Statement: Due to lack of the relevant secondary data, the Machine learning model proposed in this study was learned based on the domain expert opinions and the narrative of relevant literature, such as the English National Study of Flooding and Health (NSFH, 2020) available at <https://bit.ly/3eXiKwt>.

Conflicts of Interest: The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Abbreviations

The following abbreviations are used in this manuscript:

BN	Bayesian Network
CI	Conditional Probability
CQALY	Change in QALY
CU	Change in Utility
DAG	Directed Acyclic Graph
EA	Environment Agency
EWS	Early Warning System
LOSI	Loss Of Sentimental Items
Lsever	Less severe depression
Msever	More severe depression
NSFH	English National Study of Flooding and Health
PHE	Public Health England
PPD	Prevalence of Probable Depression
PTSD	Post-Traumatic Stress Disorder
QALY	Quality-Adjusted Life Year
WTP	Willingness to Pay

References

1. Munich, R. (2014). *Topics annual review: Natural disasters 2014* Munchener Rckversicherungs-Gesellschaft.
2. UNISDR (2011). *Global assessment report on disaster individual flood protection*. Revealing risk, Redefining development.
3. World Economic Forum. (2018). *The Global Risks Report 2018 13th edition*. World Economic Forum. Geneva, Switzerland.
4. Hammond, M. J., Chen, A. S., Djordjevic, S., Butler, D., and Mark, O. (2015). Urban flood impact assessment: A state-of-the-art review. *Urban Water Journal*, **12**(1), 14-29.
5. Chatterton, J., Clarke, C., Daly, E., Dawks, S., Elding, C., Fenn, T., ... & Salado, R. (2016). The costs and impacts of the winter 2013 to 2014 floods. *Report SC140025*, 1.
6. Mechler, R. (2016). Reviewing estimates of economic efficiency of disaster risk management: opportunities & limitations of using risk-based cost-benefit analysis. *Nat. Hazs*, **81**(3), 2121-47.
7. Walker-Springett, K., Butler, C., & Adger, W. N. (2017). Wellbeing in the aftermath of floods. *Health & place*, **43**, 66-74.
8. Lamond, J.E., Joseph, R.D. and Proverbs, D.G. (2015). An exploration of factors affecting the long term psychological impact and deterioration of mental health in flooded households. *Environmental research*, **140**, 325-334.
9. Prettenthaler, F., Koberl, J., & Bird, D. N. (2016). ‘Weather Value at Risk’: A uniform approach to describe and compare sectoral income risks from climate change. *Science of the Total Environment*, **543**, 1010-1018.
10. Fewtrel, L. J., Kay, D., & Ashley, R. (2008). Flooding and Health an evaluation of the health impacts of urban pluvial flooding in the UK. In *Health impact ass. for sust. water mana.*, 121-48.
11. European Council (2007). *EU directive of the European Parliament and of the European Council on the estimation and management of flood risks 2007/60/EU*.
12. Arora, P., Boyne, D., Slater, J. J., Gupta, A., Brenner, D. R., and Druzdzal, M. J. (2019). Bayesian networks for risk prediction using real-world data: a tool for precision medicine. *Value in Health*, **22**(4), 439-445.

13. Daneshkhah, A., Parham, G., Chatrabgoun, O., and Jokar, M. (2016). Approximation multivariate distribution with pair copula using the orthonormal polynomial and Legendre multiwavelets basis functions. *Communications in Statistics-Simulation and Computation*, **45(2)**, 389-419.
14. Vepa, A., Saleem, A., Rakhshan, K., Daneshkhah, A., Sedighi, T., Shohaimi, S., ... & Chakrabarti, P. (2021). Using Machine Learning Algorithms to Develop a Clinical Decision-Making Tool for COVID-19 Inpatients. *International Journal of Environmental Research and Public Health*, **18(12)**, 6228.
15. Phifer, J.F. (1990). Psychological distress and somatic symptoms after natural disaster: differential vulnerability among older adults. *Psychology and Aging*, **5(3)**, p.412.
16. Reacher, M., McKenzie, K., Lane, C., Nichols, T., Kedge, I., Iversen, A., Hepple, P., Walter, T, Laxton, C, Simpson, J. (2004). Health impacts of flooding in Lewes: a comparison of reported gastrointestinal & other illness & mental health in flooded/non-flooded households. *Comm. Dise. & PH*, **7**, 39-46.
17. Tapsell, S.M., Penning-Rowsell, E.C., Tunstall, S.M. and Wilson, T.L. (2002). Vulnerability to flooding: health and social dimensions. *Philosophical transactions of the royal society of London. Ser A: Math, Phy. & Eng. Scis.*, **360(1796)**, 1511-25.
18. Carroll, B., Morbey, H., Balogh, R. and Araoz, G. (2009). Flooded homes, broken bonds, the meaning of home, psychological processes and their impact on psychological health in a disaster. *Health & Place*, **15(2)**, pp. 540-547.
19. Mason, V., Andrews, H., Upton, D. (2010). The psychological impact of exposure to floods. *Psyc., heal. & med.*, **15(1)**, 61.
20. Bich, H., Quang, N., ThanhHa, T., DucHanh, T. & Guha-Sapir, D. (2011). Impacts of flood on health: epidemiologic evidence from Hanoi, Vietnam. *Global health action*, **4(1)**, 6356.
21. Hu, M., Zhang, X., Li, Y., Yang, H. and Tanaka, K. (2019). Flood mitigation performance of low impact development technologies under different storms for retrofitting an urbanized area. *J. of cleaner production*, **222**, 373-380.
22. Patricia S.T.D, Filipe S.F.L, Manuel, V.C.R., Paulo, M.J, and Antonio, L.P.F. (2019). Can land cover changes mitigate large floods? A reflection based on partial least squares-path modeling. *Water*, **11(4)**, p. 684.
23. Zhong, S., Yang, L., Toloo, S., Wang, Z., Tong, S., Sun, X., Crompton, D., FitzGerald, G., Huang, C. (2018). The long-term physical & psychological health impacts of flooding: a systematic mapping. *Sci. of total environment*, **626**, 165-194.
24. Marsh, T., Kirby, C., Muchan, K., Barker, L., Henderson, E, Hannaford, J. (2016). The winter floods of 2015/2016 in the UK-a review. In *7th Int. Conf. on Hydroinformatics*.
25. Schmitt, L.H., Graham, H.M. and White, P.C. (2016). Economic evaluations of the health impacts of weather-related extreme events: a scoping review. *International journal of environmental research and public health*, **13(11)**, p.1105.
26. Matsushima, K., Onishi, M. and Kobayashi, K. (2007). The winter floods of 2015/2016 in the UK-a review. In *2007 IEEE Int. Conf. on Systems, Man and Cybernetics*, 1848-1853.
27. Navrud, S., Huu Tuan, T. and Duc Tinh, B. (2012). Estimating the welfare loss to households from natural disasters in developing countries: a contingent valuation study of flooding in Vietnam. *Global health action*, **5(1)**, p. 17609.
28. DEFRA (2004). *The appraisal of human-related intangible impacts of flooding. R&D Technical Report FD2005/TR*.
29. DEFRA (2012). *UK climate change risk assessment. Evidence Report*.
30. Crichton, D. (1999). The Risk Triangle. In: *Ingleton, J., Ed., Nat. Disaster Manag., Tudor Rose, London*, **566**, 102-103.
31. Gain, A.K., Mojtahed, V., Biscaro, C., Balbi, S. & Giupponi, C. (2015). An integrated approach of flood risk assessment in the eastern part of Dhaka City. *Nat. Hazards*, **79(3)**, 1499-1530.
32. Jonkman, S.N., Kok, M. & Vrijling, J.K. (2018). Flood risk assessment in the Netherlands: A case study for dike ring South Holland. *Risk Analysis: An Int. Jou.*, **28(5)**, pp. 1357-1374.
33. Balbi, S., Villa, F, Mojtahed, V., Hegetschweiler, K.T. and Giupponi, C. (2016). A spatial Bayesian network model to assess the benefits of early warning for urban flood risk to people. *Natural Hazards & Earth System Sciences*, **16(6)**.
34. Sibley, A. (2010). Analysis of extreme rainfall and flooding in Cumbria. *Weather*, **55(11)**, pp.287-292.
35. Coles, D., Yu, D., Wilbey, R.L., Green, D, Herring, Z. (2017). Beyond 'flood hotspots': modelling emergency service accessibility during flooding in York. *JoH*, **566**, 419-36.
36. Lowe, D., Ebi, K.L. and Forsberg, B. (2013). Factors increasing vulnerability to health effects before, during and after floods. *Int. J. of envi. res. and public health*, **10(12)**, 7015-67.
37. Jermacane, D., Waite, T.D., Beck, C.R., Bone, A., Amlôt, R., Reacher, M., Kovats, S., Armstrong, B., Leonardi, G., Rubin, G.J. & Oliver, I. (2018). The English National Cohort Study of Flooding and Health: the change in the prevalence of psychological morbidity at year two. *BMC PH*, **18(1)**, 330.
38. Huang, P., ZTan, H., Liu, A., Feng, S. and Chen, M. (2010). Prediction of posttraumatic stress disorder among adults in flood district. *BMC Public Health*, **10(1)**, p. 207.
39. Fenton, N. and Neil, M. (2012). *Risk assessment and decision analysis with Bayesian networks* Crc Press.
40. Uusitalo, L. (2007). Advantages & challenges of BNs in environmental modelling. *Ecol. modelling*, **203(3-4)**, 312-318.
41. Sedighi, T., and Varga, L. (2019). CECAN Evaluation and Policy Practice Note (EPPN) for policy analysts and evaluators-A Bayesian Network for Policy Evaluation.
42. Barton, D.N., Kuikka, S., Varis, O., Uusitalo, L., Henriksen, H.J., Borsuk, M., de la Hera, A., Farmani, R., Johnson, S. & Linnell, J.D. (2011). BNs in environmental and resource management. *Integ. envi. assessment and management*, **8(3)**, 418-29.
43. Bedford, T., Wilson, K. J., and Daneshkhah, A. (2014). Assessing parameter uncertainty on coupled models using minimum information methods. *Reliability Engineering & System Safety*, **125**, 3-12.

44. Jensen, K. L., Toftum, J., & Friis-Hansen, P. (2009). A Bayesian Network approach to the evaluation of building design and its consequences for employee performance and operational costs. *Building and Environment*, 44(3), 456–462.
45. Sedighi, T., and Varga, L. (2021). Evaluating the Bovine Tuberculosis Eradication Mechanism and Its Risk Factors in England's Cattle Farms. *International Journal of Environmental Research and Public Health*, 18(7), 3451.
46. Chatrabgoun, O., Hosseinian-Far, A., Chang, V., Stocks, N. G., and Daneshkhah, A. (2018). Approximating non-Gaussian Bayesian networks using minimum information vine model with applications in financial modelling. *Journal of computational science*, 24, 266-276.
47. Nunes, J., Barbosa, M., Silva, L., Gorgonio, K., Almeida, H., and Perkusich, A. (2018). Issues in the probability elicitation process of expert-based Bayesian networks. *Enhanced Expert Systems*, No. 5.
48. Zhou, Y., Fenton, N., and Neil, M. (2014). Bayesian network approach to multinomial parameter learning using data and expert judgments. *International Journal of Approximate Reasoning*, 55(5), 1252-1268.
49. Lee, C., and van Beek, P. (2017, May). Metaheuristics for score-and-search Bayesian network structure learning. In *Canadian Conference on Artificial Intelligence* (pp. 129-141). Springer, Cham.
50. Sedighi, T., (2019). Using Dynamic and Hybrid Bayesian Network for Policy Decision Making. *International Journal of Strategic Engineering (IJoSE)*, 2(2), p.22-34.
51. Smith, J. Q., and Daneshkhah, A. (2010). On the robustness of Bayesian networks to learning from non-conjugate sampling. *International Journal of Approximate Reasoning*, 51(5), 558-572.
52. Daneshkhah, A., and Oakley, J. E. (2010). Eliciting multivariate probability distributions. *Rethinking risk measurement and reporting*, 1, 23.
53. Waite, T. D., Chaintarli, K., Beck, C. R., Bone, A., Amlôt, R., Kovats, S., Reacher, M., Armstrong, B., Leonardi, G. and Rubin, G. J. (2017). The English national cohort study of flooding and health: cross-sectional analysis of mental health outcomes at year one. *BMC public health*, 17, 1-9.
54. National Institute for Clinical Excellence. (2018). *Depression in adults: treatment and management*.
55. Oakley, J. E. (2009). Decision-theoretic sensitivity analysis for complex computer models. *Technometrics*, 51(2), 121-129.
56. Daneshkhah, A., Hosseinian-Far, A., and Chatrabgoun, O. (2017). Sustainable maintenance strategy under uncertainty in the lifetime distribution of deteriorating assets. In *Strategic Engineering for Cloud Computing and Big Data Analytics* (pp. 29-50). Springer, Cham.
57. McCabe, C., Claxton, K., & Culyer, A. J. (2008). The NICE cost-effectiveness threshold. *Pharmacoeconomics*, 26(9), 733-744.
58. Vallejo-Torres, L., García-Lorenzo, B., Castilla, I., Valcárcel-Nazco, C., García-Pérez, L., Linertová, R., ... & Serrano-Aguilar, P. (2016). On the estimation of the cost-effectiveness threshold: why, what, how?. *Value in Health*, 19(5), 558-566.
59. Rounsevell, M.D.A., Metzger, M.J. (2010). Developing qualitative scenario storylines for environmental change assessment. *Wiley Inter. Reviews: Climate Change*, 1(4), 606–619.